

**ANNEX I**

**SUMMARY OF PRODUCT CHARACTERISTICS**

## 1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Avishield IBD INT lyophilisate for oculonasal suspension/use in drinking water for chickens

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each dose contains:

### Active substance:

Infectious bursal disease virus, IM strain VMG 91, Live

$10^{4.0}$  to  $10^{5.0}$  TCID<sub>50</sub>\*

\*TCID<sub>50</sub> = 50% tissue culture infective dose.

### Excipients:

Qualitative composition of excipients and other constituents
Povidone K 25
Bacto peptone
Monosodium glutamate
Potassium dihydrogen phosphate
Potassium hydroxide

Cream to reddish coloured lyophilisate.

## 3. CLINICAL INFORMATION

### 3.1 Target species

Chickens.

### 3.2 Indications for use for each target species

For active immunisation of chickens (broilers, future layers and breeders), with maternally derived antibodies, to prevent mortality and clinical disease, due to infection caused by infectious bursal disease viruses.

Onset of immunity: 2 weeks after vaccination.

Duration of immunity: 4 weeks after vaccination.

### 3.3 Contraindications

None.

### 3.4 Special warnings

Please refer to section 'Administration routes and dosage'.

Vaccinate healthy animals only.

### 3.5 Special precautions for use

Special precautions for safe use in the target species:

The vaccine strain can spread to susceptible, unvaccinated chickens for at least 10 days following vaccination. The vaccine virus has shown the potential to increase in virulence on bird to bird passage and may cause immunosuppression but does not induce clinical signs of disease. It is very important to take measures to ensure that the vaccine strain does not spread to unvaccinated chickens.

It is possible that the vaccine viruses can be spread to susceptible non-target species. Care should be taken to ensure that the vaccine virus does not spread to unvaccinated birds. Therefore, all birds in a flock should be vaccinated at the same time to reduce the risk of bird to bird transmission. Vaccinated birds should not be mixed with unvaccinated birds. Hygiene measures should be taken to prevent spread to other flocks. Vaccination of all chickens on the premises is recommended.

Housing needs to be disinfected prior to restocking.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Wash and disinfect hands and equipment after vaccination.

Special precautions for the protection of the environment:

Not applicable.

### 3.6 Adverse events

Chickens.

Very common (>1 animal / 10 animals treated):	Bursa of Fabricius lymphocyte depletion <sup>a</sup>
--	--

<sup>a</sup>Mild to moderate; 7 days after vaccine take (bursal lesion score 2.4). This depletion decreases and is followed by lymphocyte repopulation and complete regeneration of the bursa by day 28 post vaccination (bursal lesion score 0.2).

Reporting adverse events is important. It allows continuous safety monitoring of a veterinary medicinal product. Reports should be sent, preferably via a veterinarian, to either the marketing authorisation holder or its local representative or the national competent authority via the national reporting system. See the package leaflet for respective contact details.

### 3.7 Use during pregnancy, lactation or lay

Laying birds:

Do not use in birds in lay or within 4 weeks before the start of the laying period.

### 3.8 Interaction with other medicinal products and other forms of interaction

No information is available on the safety and efficacy of this vaccine when used with any other veterinary medicinal product. A decision to use this vaccine before or after any other veterinary medicinal product therefore needs to be made on a case by case basis.

### 3.9 Administration routes and dosage

One dose of vaccine should be administered to each chicken in drinking water, or by the oculonasal route from 8 days of age depending on the level of maternally derived antibodies (MDA) in the flock. Lyophilisate reconstituted in 100 mL of water appears as non-transparent suspension with low precipitation.

The optimal vaccination date is influenced by a number of factors, such as status of MDA, type of bird, infection pressure, housing and management conditions.

MDA can interfere with the immunity induced by live infectious bursal disease (IBD) vaccines, so the optimum age for vaccination depends on both the level of residual MDA against IBD virus in the flock and the ability of the vaccine strain to induce the required level of immunity in the presence of MDA. To predict the age when the MDA titre has sufficiently decreased to allow effective vaccination (“break-through titre”), testing of serum samples of at least 18 chicks by serology and application of the “Deventer formula” is advised. A break-through titre of 125 should be used.

The Deventer formula is as follows:

Vaccination age = { (log<sub>2</sub> titre bird% - log<sub>2</sub> breakthrough) x t<sub>1/2</sub> } + age at sampling + correction 0-4

In which:

- bird% = ELISA titre of the bird representing a certain percentage of the flock
- breakthrough = breakthrough (ELISA) titre of the vaccine to be used
- t<sub>1/2</sub> = half-life time (ELISA) of the antibodies in the type of chickens being sampled
- age at sampling = age of the birds at sampling
- correction 0-4 = extra days when the sampling was done at 0 to 4 days of age.

A high homogeneity of MDA levels in the flock is important to define the correct timing of vaccination and guarantees a better active immune response to the vaccine. In case of a non-homogenous flock where antibody levels vary widely between birds (i.e. CV greater than 30%), or the stock originates from different sources, it is recommended to repeat the vaccination.

In such cases, timing of the first and second vaccination should be determined simultaneously, with two different percentages (corresponding to the percentages of the flock that can be efficaciously vaccinated) of all serum samples collected on the sampling day, using the Deventer formula.

### 1. In drinking water use

- Reconstitute the vaccine in a small amount of cool and clean water without traces of chlorine, other disinfectants or impurities, in a number of doses corresponding to the number of birds to be vaccinated. Where the number of birds is between the standard dosages, the next higher dosage should be used.
- The vaccine should be reconstituted immediately before use.
- Measure the correct volume of water for the number of birds to be vaccinated. The volume of water for dilution depends on the age of the birds, breed, housing conditions and weather conditions.
- The reconstituted vaccine should be diluted in the amount of water which will be consumed within 1.5 to 2.0 hours (taking into account the different types of drinking systems for poultry).
- In order to determine the quantity of water in which the vaccine will be diluted, measure the volume of water consumed within a two hours period one day before vaccination.
- As a guideline for younger chickens (until 3rd week of life), apply the reconstituted vaccine to cold and fresh water at the rate of 1 000 doses of vaccine to 1 litre of water per day of age for 1 000 chickens, e.g. 8 litres would be needed for 1 000, 8 day old chickens.
- In order to make the birds thirsty, withdraw the supply of drinking water up to 2 hours prior to immunisation (birds drinking behaviour varies, depending on the air temperature, type of birds, breed, management, weather conditions).
- The drinking system should be clean, without traces of chlorine, other disinfectants or impurities.
- If needed, turn the lights down low when the water is turned off. After the vaccine is in the drinking system, increase light intensity again. Increased light intensity will stimulate the birds to look for food and water.
- Always make sure that there is food available when vaccinating. Birds will not drink if they have no food to eat.

### 2. Oculonasal use

- Reconstitute 1 000 doses of the vaccine in 100 ml distilled water.

- A dose of reconstituted vaccine is 0.1 ml, i.e. two drops, irrespective of poultry age, weight and type. Instill one drop into an eye and one drop into a nostril.

### **3.10 Symptoms of overdose (and where applicable, emergency procedures and antidotes)**

After the administration of a 10-fold overdose, no adverse reactions other than those described in section Adverse events were observed.

### **3.11 Special restrictions for use and special conditions for use, including restrictions on the use of antimicrobial and antiparasitic veterinary medicinal products in order to limit the risk of development of resistance**

Not applicable.

### **3.12 Withdrawal periods**

Zero days.

## **4. IMMUNOLOGICAL INFORMATION**

### **4.1 ATCvet code: QI01AD09**

To stimulate active immunity against infectious bursal disease virus in chickens.

The vaccine strain is an intermediate strain with an average bursal lesion score of 0.2 at 28 days after administration of 10-times the maximum dose.

## **5. PHARMACEUTICAL PARTICULARS**

### **5.1 Major incompatibilities**

Do not mix with any other veterinary medicinal product.

### **5.2 Shelf life**

Shelf life of the veterinary medicinal product as packaged for sale: 2 years

Shelf life after reconstitution according to directions: 3 hours.

### **5.3 Special precautions for storage**

Store in a refrigerator (2 °C – 8 °C).

Do not freeze.

Protect from light.

### **5.4 Nature and composition of immediate packaging**

The vaccine is filled into 4 ml or 10 ml colourless glass vials (type I), which are closed with brombutyl rubber stoppers and sealed with aluminium caps.

Pack sizes:

Carton box with 10 vials of 1 000 doses of vaccine.

Carton box with 10 vials of 2 500 doses of vaccine.

Carton box with 10 vials of 5 000 doses of vaccine.

Not all pack sizes may be marketed.

**5.5 Special precautions for the disposal of unused veterinary medicinal products or waste materials derived from the use of such products**

Medicines should not be disposed of via wastewater or household waste.

Use take-back schemes for the disposal of any unused veterinary medicinal product or waste materials derived thereof in accordance with local requirements and with any national collection systems applicable to the veterinary medicinal product concerned.

**6. NAME OF THE MARKETING AUTHORISATION HOLDER**

Genera d.d..

**7. MARKETING AUTHORISATION NUMBER(S)**

VPA/0405/004/001

**8. DATE OF FIRST AUTHORISATION**

Date of first authorisation: 07/01/2019

**9. DATE OF THE LAST REVISION OF THE SUMMARY OF THE PRODUCT CHARACTERISTICS**

<{DD/MM/YYYY}>

**10. CLASSIFICATION OF VETERINARY MEDICINAL PRODUCTS**

Veterinary medicinal product subject to prescription.

Detailed information on this veterinary medicinal product is available in the [Union Product Database \(https://medicines.health.europa.eu/veterinary\)](https://medicines.health.europa.eu/veterinary).

